

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Editors: W. Clegg and D. G. Watson

**Bis[ $\mu$ -3-(*N'*-methyl-4,4'-bipyridinium-1-yl)propionamidato]bis[*cis*-diammineplatinum(II)] hexaperchlorate dihydrate: a head-to-tail isomer**

**Ken Sakai, Yoshimi Yokoyama, Hideki Hama, Kaoru Kato, Yoshie Ikuta, Taro Tsubomura and Tomoaki Tanase**

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site provided that this cover page is retained. Republication of this article or its storage in electronic databases or the like is not permitted without prior permission in writing from the IUCr.

Ken Sakai,<sup>a\*</sup> Yoshimi Yokoyama,<sup>a</sup> Hideki Hama,<sup>a</sup> Kaoru Kato,<sup>a</sup> Yoshie Ikuta,<sup>b</sup> Taro Tsubomura<sup>b</sup> and Tomoaki Tanase<sup>c</sup>

<sup>a</sup>Department of Applied Chemistry, Faculty of Science, Tokyo University of Science, Kagurazaka 1-3, Shinjuku-ku, Tokyo 162-8601, Japan, <sup>b</sup>Department of Applied Chemistry, Faculty of Engineering, Seikei University, Kichijoji-Kitamachi 3-3-1, Musashino, Tokyo 180-8633, Japan, and <sup>c</sup>Department of Chemistry, Faculty of Science, Nara Women's University, Kitauoya-higashi-machi, Nara 630-8285, Japan

Correspondence e-mail: ksakai@rs.kagu.tus.ac.jp

#### Key indicators

Single-crystal X-ray study  
 $T = 296$  K  
 Mean  $\sigma(\text{C}-\text{C}) = 0.021$  Å  
 Some non-H atoms missing  
 Disorder in solvent or counterion  
 $R$  factor = 0.060  
 $wR$  factor = 0.121  
 Data-to-parameter ratio = 18.5

For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.

## Bis[ $\mu$ -3-( $N'$ -methyl-4,4'-bipyridinium-1-yl)propionamidato]bis[*cis*-diammineplatinum(II)] hexaperchlorate dihydrate: a head-to-tail isomer


In the title compound,  $[\text{Pt}_2(\mu\text{-C}_{14}\text{H}_{16}\text{N}_3\text{O})_2(\text{NH}_3)_4](\text{ClO}_4)_6 \cdot 2\text{H}_2\text{O}$ , the diplatinum(II) cation is found to be a head-to-tail isomer. The intradimer Pt—Pt distance [3.0304 (7) Å] is much shorter than the value of 3.0852 (13) Å reported for the analogous *cis*-diammineplatinum(II) dimer bridged by 2-( $N'$ -methyl-4,4'-bipyridinium-1-yl)acetamidates [Sakai, Ikuta, Tsubomura, Kato, Yokoyama, Kajiwara & Ito (2003). *Acta Cryst. E* **59**, m780–m783], showing that the electron density at the metal centers is higher in the title system than in the previously reported system. The shortest interdimer Pt...Pt distance is 8.4585 (8) Å.

#### Comment


We previously reported that amidate-bridged *cis*-diammineplatinum(II) dimers, *viz.*  $[\text{Pt}_2(\text{NH}_3)_4(\mu\text{-amidato})_2]^{2+}$  (amidate is acetamidate,  $\alpha$ -pyrrolidinonate,  $\alpha$ -pyridonate *etc.*), are generally active as  $\text{H}_2$ -evolving catalysts in a well known photosystem consisting of edta (usually, ethylenediaminetetraacetic acid disodium salt),  $\text{Ru}(\text{bpy})_3^{2+}$  (bpy is 2,2'-bipyridine), and methylviologen (usually,  $N,N'$ -dimethyl-4,4'-bipyridinium dichloride) (Sakai & Matsumoto, 1990; Sakai *et al.*, 1993). It should be noted here that two geometrical isomers, head-to-head (HH) and head-to-tail (HT), are possible for this class of doubly-bridged dimers, because of the asymmetric feature of amidate N—C—O units. In addition, edta,  $\text{Ru}(\text{bpy})_3^{2+}$  and methylviologen, respectively, serve as a sacrificial electron donor, a photosensitizer and an electron relay. One of our interests over many years has concentrated on the development of molecular devices which replace three major components, that is,  $\text{Ru}(\text{bpy})_3^{2+}$ , methylviologen and the  $\text{Pt}^{\text{II}}_2$  dimer. Such devices are considered as one-component systems. Considerable attention has also been paid to the studies on two-component systems in which the photochemical reduction of water into  $\text{H}_2$  by edta is driven by  $\text{Ru}(\text{bpy})_3^{2+}$  and a hybrid compound made up of both the methylviologen and the  $\text{H}_2$ -evolving  $\text{Pt}^{\text{II}}_2$  units. In this context, the title compound, HT- $[\text{Pt}_2(\text{NH}_3)_4(\mu\text{-L1})_2](\text{ClO}_4)_6 \cdot 2\text{H}_2\text{O}$  [(I); L1 is 3-( $N'$ -methyl-4,4'-bipyridinium-1-yl)propionamidate], together with an analogous dimer, HT- $[\text{Pt}_2(\text{NH}_3)_4(\mu\text{-L2})_2](\text{ClO}_4)_6 \cdot 2\text{H}_2\text{O}$  [(II); Sakai, Ikuta *et al.*, 2003; L2 is 2-( $N'$ -methyl-4,4'-bipyridinium-1-yl)acetamidate], were first prepared in 1994 (Ikuta *et al.*, 1994), and were found to be somewhat effective. It has been confirmed that the  $\text{H}_2$ -evolving activity of (I), evaluated under the three-component system, is slightly higher than that of the acetamidate-bridged analog (Sakai *et al.*, 1993), and is *ca.* 1.5 times higher than that of (II) (Sakai *et al.*, 1995). Importantly, preliminary experiments have shown that the photolysis of a solution containing edta,  $\text{Ru}(\text{bpy})_3^{2+}$ , and either (I) or (II), in the absence of

Received 5 April 2004  
 Accepted 20 April 2004  
 Online 30 April 2004


methylviologen, also results in evolution of H<sub>2</sub> in a relatively high quantum efficiency (Sakai *et al.*, 1995). Nevertheless, it took a long time to refine the reliable synthetic methods yielding (I), (II) and the related complexes (Sakai, Shiomi *et al.*, 2003; Sakai, Ikuta *et al.*, 2003). We now report the synthesis and crystal structure of (I).



A dinuclear Pt<sup>II</sup><sub>2</sub> cation, six perchlorate anions and apparently 1.5 molecules of water are found in the asymmetric unit of (I). As previously reported for several analogous dimers doubly bridged by chain amidates, the binding directions of two bridging amidates could be rationally determined (see *Experimental*). As a result, the complex has been shown to be a head-to-tail isomer in the crystal structure, even though the



**Figure 1**  
The structure of (I), showing the atom-labeling scheme. Displacement ellipsoids are drawn at the 50% probability level. Perchlorate ions and water molecules have been omitted for clarity.



**Figure 2**

A stereoview for the crystal packing of (I), with atoms represented by spheres of arbitrary sizes. The minor disorder components of the perchlorate ions have been omitted for clarity.

complex is likely to give a mixture of both the HH and the HT isomer in solution, as previously observed for the related amidate-bridged Pt<sup>II</sup><sub>2</sub> dimers (see, for example, Sakai, Tanaka *et al.*, 1998).

The intradimer Pt—Pt distance in (I) [3.0304 (7) Å] is much shorter than that reported for (II) [3.0852 (13) Å] (Sakai, Ikuta *et al.*, 2003). The considerably long Pt<sup>II</sup>—Pt<sup>II</sup> distance in (II) was previously interpreted in terms of the fact that the positively charged bipyridinium moieties in (II) serve as electron-withdrawing groups to diminish the electron density at the Pt<sup>II</sup> centers, leading to the weakening of the ‘mutual dative bonds’ within the diplatinum cation [two Pt<sup>II</sup> centers in close proximity are considered to form a dative Pt(5d<sub>z<sup>2</sup>) → Pt(6p or 6s) bond with one another; Connick *et al.*, 1997]. The shorter Pt—Pt distance in (I) is consistent with the fact that the bipyridinium moieties in (I) serve as poorer electron-withdrawing groups, because the distance between the bipyridinium unit and the Pt<sup>II</sup><sub>2</sub> core in (I) is longer than that in (II). The N(bipyridinium)—C(bridging amidate) distances in (I) [N2—C1 = 3.831 (18) Å and N5—C15 = 3.781 (17) Å] are *ca* 1.4 Å longer than those reported for (II) [2.41 (3) and 2.46 (3) Å; Sakai, Ikuta *et al.* 2003], where the N(bipyridinium) atom denotes the N atom of a viologen unit which is closer to the diplatinum entity.</sub>

The dihedral angle between the two Pt coordination planes within the dimeric unit ( $\tau$ ) and the average torsional twist of them about the Pt—Pt axis ( $\omega$ ) are estimated as  $\tau = 37.8$  (3)° and  $\omega = 3.8$  (9)° (see also Table 1), where  $\omega = 0^\circ$  denotes that the two Pt coordination planes stack in an eclipsed fashion. The corresponding values were reported as  $\tau = 40.8$  (5)° and  $\omega = 5$  (1)° for (II) (Sakai, Ikuta *et al.*, 2003). Atoms Pt1 and Pt2

are shifted out of their coordination planes by 0.073 (5) and 0.065 (5) Å, respectively, in such a manner that they have an attractive interaction with one another.

The pyridinium plane directly attached to the propionamide unit is inclined with respect to the bridging O—C—N unit by 89.8 (7)° for N2/C4—C8 and 6.8 (24)° for N5/C18—C22. The twist angles of the two pyridinium planes within the individual bipyridinium moieties are 32.0 (7)° for the bipyridinium involving N2 and N3, and 16.2 (6)° for that involving N5 and N6. As previously discussed for (II) (Sakai, Ikuta *et al.*, 2003), these values are both effectively larger than that observed for the uncoordinated ligand L2H (a protonated form of L2) [6.1 (2)°; Sakai *et al.*, 1997], implying that part of the electron density at the Pt<sup>II</sup><sub>2</sub> core is shifted towards the π\*(bipyridinium) orbitals, giving rise to the lower aromaticity at the central C—C bonds of bipyridinium units. It should be also noted that the relatively large twist of 32.0 (7)° observed for the bipyridinium unit involving N2 and N3 is, in part, due to the strain enhanced by the intermolecular electrostatic interactions between the anions and the cations.

Fig. 2 shows a crystal packing view of (I). The shortest intermolecular Pt...Pt distance is observed as Pt1...Pt2(x, y - 1, z) = 8.4585 (8) Å (Table 1), confirming the lack of any intermolecular Pt—Pt interaction in the crystal structure. The crystal packing is stabilized both by the electrostatic interactions between the complex cations and perchlorate anions and by hydrogen bonds formed between the amines and the O atoms of perchlorates as well as those between the water molecules and the O atoms of perchlorates (Tables 2 and 3).

## Experimental

*N*-Methyl-4,4'-bipyridinium iodide was prepared according to the literature method of van Emon *et al.* (1986). *N*-(2-Carbamoyl-ethyl)-*N'*-methyl-4,4'-bipyridinium diperchlorate was prepared in the same manner as previously described for the synthesis of *N*-(carbamoyl-methyl)-*N'*-methyl-4,4'-bipyridinium diperchlorate dihydrate (Sakai *et al.*, 1997). A solution of 3-bromopropionamide (10 mmol, 1.52 g) and *N*-methyl-4,4'-bipyridinium iodide (10 mmol, 2.98 g) in dry methanol (25 ml; dried over molecular sieve 4 Å) was refluxed for 48 h. After the solution was cooled to room temperature, the resulting orange powder was collected by filtration and air-dried. The product was then converted into the chloride salt using anion exchange resin (Amberlite IRA-400 in the Cl form). The chloride salt was further treated with two equivalents of silver perchlorate in water in the dark at 333 K for 1 h, followed by removal of AgCl by filtration. The filtrate was evaporated at 313–323 K almost to dryness. (Caution: perchlorate salts are potentially explosive and must be handled with care, even though we never experienced any difficulty in handling this material.) To the oily solution was added an excess of ethanol to give the product as a colorless powder which was recrystallized from a water–ethanol mixture (1:7 v/v) (yield 72%). Analysis calculated for C<sub>14</sub>H<sub>17</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>9</sub>: C 38.03, H 3.87, N 9.50%; found: C 38.09, H 3.83, N 9.34%. <sup>1</sup>H NMR (D<sub>2</sub>O, 296 K): δ 3.17 (*t*, 2H, *J* = 6.3 Hz), 4.51 (*s*, 3H), 5.02 (*t*, 2H, *J* = 6.3 Hz), 8.53 (*d*, 2H, *J* = 7.1 Hz), 8.56 (*d*, 2H, *J* = 6.8 Hz), 9.07 (*d*, 2H, *J* = 6.6 Hz), 9.17 (*d*, 2H, *J* = 7.1 Hz).

Compound (I) was prepared as follows: to an aqueous solution of *cis*-[Pt(NH<sub>3</sub>)<sub>2</sub>(OH<sub>2</sub>)<sub>2</sub>](ClO<sub>4</sub>)<sub>2</sub> (0.2 mmol Pt/2 ml H<sub>2</sub>O), prepared as

previously described (Sakai, Takeshita *et al.*, 1998), was added *N*-(2-carbamoyl-ethyl)-*N'*-methyl-4,4'-bipyridinium diperchlorate (0.32 mmol). The solution was heated at 343 K for 4 h followed by addition of a solution of NaClO<sub>4</sub> (6.0 mmol) in water (0.35 ml). After the solution was filtered for the removal of insoluble materials, the filtrate was left to stand at room temperature for a few days. The yellow needles deposited were collected by filtration and air-dried (yield 15%). Analysis calculated for C<sub>28</sub>H<sub>48</sub>Cl<sub>6</sub>N<sub>10</sub>O<sub>28</sub>Pt<sub>2</sub>: C 21.34, H 3.07, N 8.89%; found: C 21.47, H 2.60, N 8.55%. <sup>1</sup>H NMR (D<sub>2</sub>O, 296 K): δ 3.10–3.16 (*m*, 2H), 4.50 (*s*, 3H), 8.51–8.57 (*m*, 4H), 9.05–9.16 (*m*, 4H), where the multiplet (2H) derived from the methylene unit attached to the bipyridinium unit overlapped the solvent signal (4.8–4.9 p.p.m.). Although only 1.5 molecules of water could be located in the formula unit, we tentatively assumed that this compound is a dihydrate product, and this is reflected in the *Crystal data* as presented. The crystals of (I) are stable in air at room temperature. A diffraction-quality single crystal was mounted on a glass fibre with epoxy resin.

### Crystal data

[Pt <sub>2</sub> (C <sub>14</sub> H <sub>16</sub> N <sub>3</sub> O) <sub>2</sub> (NH <sub>3</sub> ) <sub>4</sub> ]- (ClO <sub>4</sub> ) <sub>6</sub> ·2H <sub>2</sub> O	<i>Z</i> = 2
<i>M<sub>r</sub></i> = 1575.64	<i>D<sub>x</sub></i> = 2.049 Mg m <sup>-3</sup>
Triclinic, <i>P1</i>	Mo <i>K</i> α radiation
<i>a</i> = 10.0927 (5) Å	Cell parameters from 2521 reflections
<i>b</i> = 11.0827 (5) Å	<i>θ</i> = 1.9–27.5°
<i>c</i> = 24.4798 (10) Å	<i>μ</i> = 5.88 mm <sup>-1</sup>
<i>α</i> = 78.116 (2)°	<i>T</i> = 296 (2) K
<i>β</i> = 79.305 (2)°	Needle, yellow
<i>γ</i> = 74.210 (2)°	0.12 × 0.04 × 0.03 mm
<i>V</i> = 2553.9 (2) Å <sup>3</sup>	

### Data collection

Bruker SMART APEX CCD-detector diffractometer	11 715 independent reflections
<i>ω</i> scans	3519 reflections with <i>I</i> > 2σ( <i>I</i> )
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	<i>R</i> <sub>int</sub> = 0.122
<i>T</i> <sub>min</sub> = 0.666, <i>T</i> <sub>max</sub> = 0.831	<i>θ</i> <sub>max</sub> = 27.5°
24 579 measured reflections	<i>h</i> = -12 → 13
	<i>k</i> = -13 → 14
	<i>l</i> = -25 → 31

### Refinement

Refinement on <i>F</i> <sup>2</sup>	H-atom parameters constrained
<i>R</i> [ <i>F</i> <sup>2</sup> > 2σ( <i>F</i> <sup>2</sup> )] = 0.060	<i>w</i> = 1/[σ <sup>2</sup> ( <i>F<sub>o</sub></i> <sup>2</sup> ) + (0.0288 <i>P</i> ) <sup>2</sup> ]
<i>wR</i> ( <i>F</i> <sup>2</sup> ) = 0.121	where <i>P</i> = ( <i>F<sub>o</sub></i> <sup>2</sup> + 2 <i>F<sub>c</sub></i> <sup>2</sup> )/3
<i>S</i> = 0.79	(Δ/ <i>σ</i> ) <sub>max</sub> < 0.001
11715 reflections	Δ <i>ρ</i> <sub>max</sub> = 0.82 e Å <sup>-3</sup>
633 parameters	Δ <i>ρ</i> <sub>min</sub> = -0.82 e Å <sup>-3</sup>

**Table 1**

Selected geometric parameters (Å, °).

Pt1—O2	2.041 (8)	Pt2—N4	1.985 (9)
Pt1—N1	2.007 (11)	Pt2—N7	2.009 (10)
Pt1—N9	2.038 (10)	Pt2—N8	2.057 (8)
Pt1—N10	2.021 (9)	Pt1—Pt2	3.0304 (7)
Pt2—O1	1.997 (9)	Pt1—Pt2 <sup>i</sup>	8.4585 (8)
N1—Pt1—N10	88.3 (4)	N4—Pt2—O1	92.1 (4)
N10—Pt1—N9	90.2 (4)	N4—Pt2—N7	87.6 (4)
N1—Pt1—O2	92.5 (4)	O1—Pt2—N8	87.5 (4)
N9—Pt1—O2	88.8 (4)	N7—Pt2—N8	92.6 (4)
N1—Pt1—N9	177.4 (4)	O1—Pt2—N7	175.9 (4)
N10—Pt1—O2	173.9 (3)	N4—Pt2—N8	176.7 (4)
O2—Pt1—Pt2—N4	-3.8 (4)	N9—Pt1—Pt2—N7	-5.0 (4)
N1—Pt1—Pt2—O1	-3.0 (4)	N10—Pt1—Pt2—N8	-3.3 (4)

Symmetry code: (i) *x*, *y* - 1, *z*.

**Table 2**  
Contact distances (Å).

O8B...O28	3.17 (3)	O27...O3 <sup>ii</sup>	3.10 (6)
O22A...O27	2.74 (4)	O28...O7B <sup>iii</sup>	3.07 (3)
O22B...O27	3.17 (5)	O28...O7A <sup>iii</sup>	3.19 (5)

Symmetry codes: (ii) 1 - x, -y, 1 - z; (iii) 1 - x, 1 - y, -z.

**Table 3**  
Hydrogen-bonding geometry (Å, °).

D—H...A	D—H	H...A	D...A	D—H...A
N7—H7C...O13B <sup>iv</sup>	0.89	2.64	2.94 (3)	101
N8—H8C...O26 <sup>v</sup>	0.89	2.28	3.159 (15)	170
N8—H8A...O21A <sup>vi</sup>	0.89	2.06	2.92 (2)	161
N8—H8B...O13A <sup>iv</sup>	0.89	2.48	3.003 (18)	118
N8—H8B...O16B <sup>iv</sup>	0.89	2.20	3.01 (2)	150
N8—H8B...O18A <sup>iv</sup>	0.89	2.34	3.155 (15)	152
N9—H9A...O22A <sup>ii</sup>	0.89	2.29	3.155 (18)	164
N9—H9C...O13B <sup>iv</sup>	0.89	2.42	3.22 (3)	150
N9—H9C...O14A <sup>iv</sup>	0.89	2.49	3.148 (18)	132
N10—H10A...O18B <sup>ii</sup>	0.89	2.49	3.00 (3)	117
N10—H10A...O20A <sup>ii</sup>	0.89	2.31	2.857 (18)	120
N10—H10A...O20B <sup>ii</sup>	0.89	2.30	3.04 (3)	141
N10—H10C...O18A <sup>iv</sup>	0.89	2.36	3.155 (15)	149
N10—H10C...O18B <sup>iv</sup>	0.89	2.28	3.00 (3)	137
N4—H4A...O9A	0.86	2.20	2.96 (3)	147
N7—H7B...O10B	0.89	2.22	3.09 (2)	165
N7—H7C...O25 <sup>v</sup>	0.89	2.59	3.151 (14)	122
N7—H7C...O26 <sup>v</sup>	0.89	2.15	3.042 (14)	177

Symmetry codes: (ii) 1 - x, -y, 1 - z; (iv) x - 1, y, z; (v) -x, 1 - y, 1 - z; (vi) 1 - x, 1 - y, 1 - z.

As recently reported for several different dimers doubly bridged by chain amidate ligands (Sakai, Kurashima *et al.*, 2003; Sakai, Ikuta *et al.*, 2003; Sakai, Shiomi *et al.*, 2003), the binding directions of O and NH for the amidate ligands were rationally determined by the results of least-squares calculations performed for two possible arrangements for each ligand as follows. A wrong combination gave an asymmetric feature with regard to the equivalent displacement parameters of the assumed O and N atoms, while an appropriate combination gave a moderate balance in these values. For O1 and N1,  $U_{\text{eq}}(\text{O1}) = 0.072$  (3) Å<sup>2</sup> and  $U_{\text{eq}}(\text{N1}) = 0.060$  (3) Å<sup>2</sup> were judged to be correct, while the reverse selection gave values of  $U_{\text{eq}}(\text{N}$  instead of O1) = 0.047 (3) Å<sup>2</sup> and  $U_{\text{eq}}(\text{O}$  instead of N1) = 0.091 (3) Å<sup>2</sup>. For O2 and N4,  $U_{\text{eq}}(\text{O2}) = 0.056$  (3) Å<sup>2</sup> and  $U_{\text{eq}}(\text{N4}) = 0.058$  (3) Å<sup>2</sup> were judged to be correct, while the reverse selection gave values of  $U_{\text{eq}}(\text{N}$  instead of O2) = 0.034 (3) Å<sup>2</sup> and  $U_{\text{eq}}(\text{O}$  instead of N4) = 0.089 (3) Å<sup>2</sup>.

Four of the six ClO<sub>4</sub><sup>-</sup> anions show orientational disorder. Around each Cl atom there are two sets of possible positions as follows: O7A/O8A/O9A/O10A and O7B/O8B/O9B/O10B around Cl2; O11A/O12A/O13A/O14A and O11B/O12B/O13B/O14B around Cl3; O15A/O16A/O17A/O18A and O15B/O16B/O17B/O18B around Cl4; and O19A/O20A/O21A/O22A and O19B/O20B/O21B/O22B around Cl5. It was assumed that the disordered O atoms around each Cl atom have the same isotropic displacement parameter. Furthermore, Cl—O distances were restrained at 1.43 (3) Å and the six O...O distances within each perchlorate anion were restrained as equal. The occu-

pancy factors of sites A and B ( $\text{sof}_A$  and  $\text{sof}_B$ ) around each Cl converged to the following values:  $\text{sof}_A(\text{Cl2}) = 0.33$  (2) and  $\text{sof}_B(\text{Cl2}) = 0.67$  (2);  $\text{sof}_A(\text{Cl3}) = 0.685$  (9) and  $\text{sof}_B(\text{Cl3}) = 0.315$  (9);  $\text{sof}_A(\text{Cl4}) = 0.702$  (8) and  $\text{sof}_B(\text{Cl4}) = 0.298$  (8); and  $\text{sof}_A(\text{Cl5}) = 0.64$  (2) and  $\text{sof}_B(\text{Cl5}) = 0.36$  (2).

All H atoms, except those of water molecules, were placed at idealized positions [C—H(methyl) = 0.96 Å, C—H(aromatic) = 0.93 Å and N—H(amine) = 0.89 Å] and included in the refinement in the riding-model approximation, with  $U_{\text{iso}}(\text{methyl H}) = 1.5U_{\text{eq}}(\text{bonded C})$ ,  $U_{\text{iso}}(\text{aromatic H}) = 1.2U_{\text{eq}}(\text{bonded C})$  and  $U_{\text{iso}}(\text{amine H}) = 1.5U_{\text{eq}}(\text{bonded N})$ . The highest peak was located 0.39 Å from atom O11A, while the deepest hole was located 0.45 Å from Pt1. Water H atoms were not located.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINTE* (Bruker, 2001); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *KENX* (Sakai, 2002); software used to prepare material for publication: *SHELXL97*, *TEXSAN* (Molecular Structure Corporation, 2001), *KENX* and *ORTEPII* (Johnson, 1976).

This work was partly supported by a Grant-in-Aid for Scientific Research (B) (No. 14340223) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

## References

- Bruker (2001). *SAINTE* (Version 6.22) and *SMART* (Version 5.625). Bruker AXS Inc., Madison, Wisconsin, USA.
- Connick, W. B., Marsh, R. E., Schaefer, W. P. & Gray, H. B. (1997). *Inorg. Chem.* **36**, 913–922.
- Emon, J. van, Hammock, B. & Seiber, J. N. (1986). *Anal. Chem.* **58**, 1866–1873.
- Ikuta, Y., Shiomi, M., Tamane, T., Tsubomura, T. & Sakai, K. (1994). *44th Symposium on Coordination Chemistry of Japan*, paper No. 1P54.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Molecular Structure Corporation (2001). *TEXSAN*. Version 1.11r1. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
- Sakai, K. (2002). *KENX*. Graphical User Interface for *SHELXL97*. Tokyo University of Science, Japan.
- Sakai, K., Ikuta, Y., Shiomi, M., Tamane, T., Tomita, Y., Tsubomura, T. & Nemoto, N. (1997). *Acta Cryst.* **C53**, 331–334.
- Sakai, K., Ikuta, Y., Tsubomura, T., Kato, K., Yokoyama, Y., Kajiwara, T. & Ito, T. (2003). *Acta Cryst.* **E59**, m780–m783.
- Sakai, K., Kizaki, Y., Tsubomura, T. & Matsumoto, K. (1993). *J. Mol. Catal.* **79**, 142–152.
- Sakai, K., Kurashima, M., Akiyama, N., Satoh, N., Kajiwara, T. & Ito, T. (2003). *Acta Cryst.* **C59**, m345–m349.
- Sakai, K. & Matsumoto, K. (1990). *J. Mol. Catal.* **62**, 1–14.
- Sakai, K., Shiomi, M., Ikuta, Y., Tomita, Y., Kitamura, Y. & Tsubomura, T. (1995). *The 1995 International Chemical Congress of Pacific Basin Societies*, paper No. INOR0518.
- Sakai, K., Shiomi, M., Tsubomura, T., Kato, K., Yokoyama, Y., Kajiwara, T. & Ito, T. (2003). *Acta Cryst.* **E59**, m559–m561.
- Sakai, K., Takeshita, M., Tanaka, Y., Ue, T., Yanagisawa, M., Kosaka, M., Tsubomura, T., Ato, M. & Nakano, T. (1998). *J. Am. Chem. Soc.* **120**, 11353–11363.
- Sakai, K., Tanaka, Y., Tsuchiya, Y., Hirata, K., Tsubomura, T., Iijima, S. & Bhattacharjee, A. (1998). *J. Am. Chem. Soc.* **120**, 8366–8379.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.